


ONEXTON[®]

(clindamycin phosphate and benzoyl peroxide)
Gel, 1.2%/3.75%

YOUR DOCTOR HAS PRESCRIBED ONEXTON— 2 MEDICATIONS IN 1 PUMP TO FIGHT ACNE

TO GET STARTED, JUST FOLLOW THESE 3 STEPS ONCE A DAY



HOW LONG DO I USE THIS?

ONEXTON has been studied up to 12 weeks, but since every person is different, **you should follow your doctor's instructions.** Do not use more ONEXTON than your doctor tells you to use.



IS TANNING OK?

Not while you are using ONEXTON...it's best to avoid tanning beds and sun lamps and to limit your time in the sun. If you are in sunlight, wear protective clothing and a broad spectrum sunscreen that is at least SPF 15.



CAN I STILL USE MAKEUP?

Yes! Once you've applied ONEXTON, you can use makeup that is noncomedogenic, which is makeup that will NOT clog your pores.



ARE THERE SIDE EFFECTS?

The most common side effect is skin irritation. If you experience severe skin irritation, stop using ONEXTON and call your doctor. This is not the only possible side effect with ONEXTON, but know that you can always talk to your doctor about any side effect that bothers you or that does not go away. **Please see the reverse side for less common, but more serious, side effects.**

INDICATION

ONEXTON (clindamycin phosphate and benzoyl peroxide) Gel, 1.2%/3.75% is a prescription medicine used on the skin (topical) to treat acne vulgaris in people 12 years and older.

Do not get ONEXTON Gel in your mouth, eyes, or vagina, or on your lips. It is not known whether ONEXTON Gel is safe and effective for use longer than 12 weeks, or in children under 12 years of age.

[Click here](#) for full Prescribing Information, including Patient Information.

YOU'VE GOT THIS!

Remember, treatment takes time to work, so it's important to apply as instructed by your doctor.

IMPORTANT SAFETY INFORMATION

Do not use ONEXTON Gel if you have: Crohn's disease, ulcerative colitis, had inflammation of the colon (colitis) or severe diarrhea with past antibiotic use. Talk with your doctor if you are not sure if you have one of these conditions.

ONEXTON Gel can cause serious side effects including:

- Inflammation of the colon (colitis). Stop using ONEXTON Gel and call your doctor right away if you have severe watery diarrhea, or bloody diarrhea.
- Allergic reactions. Stop using ONEXTON Gel, call your doctor and get help right away if you have any of the following symptoms: severe itching; swelling of your face, eyes, lips, tongue or throat; or trouble breathing.

Common side effects with ONEXTON Gel include skin irritation. Stop using ONEXTON Gel and call your doctor if you have a skin rash or your skin becomes very red, itchy or swollen. Talk to your doctor about any side effect that bothers you or that does not go away.

Tell your doctor about all the medicines and skin products you use. Especially tell your doctor if you will have surgery with general anesthesia. One of the medicines in ONEXTON Gel (clindamycin) can affect how certain medicines work when used in general anesthesia.

- ONEXTON Gel should not be used with products that contain erythromycin.
- Other skin and topical acne products may increase the irritation of your skin when used with ONEXTON Gel.

Pregnancy and Nursing Mothers: It is not known whether ONEXTON Gel will harm your unborn baby or nursing infant, so talk to your doctor if you are pregnant. If you are nursing, you and your doctor should decide if you will use ONEXTON Gel while breastfeeding.

Sun Protection: When using ONEXTON Gel, limit your time in sunlight. Avoid using tanning beds or sun lamps. If you have to be in sunlight, wear a wide-brimmed hat or other protective clothing, and a sunscreen with SPF 15 rating or higher.

You are encouraged to report negative side effects of prescription drugs. Visit www.fda.gov/medwatch or call 1-800-FDA-1088.

[Click here](#) for full Prescribing Information, including Patient Information.

YOU'RE READY TO USE ONEXTON GEL!

If you're still looking for more answers, visit www.onexton.com/FAQs

ONEXTON[®]
(clindamycin phosphate and benzoyl peroxide)
Gel, 1.2%/3.75%

ONEXTON is a trademark of Ortho Dermatologics' affiliated entities.

© All Rights Reserved. ONX.0052.USA.18